

Alcohol and Emotional Contagion: An Examination of the Spreading of Smiles in Male and
Female Drinking Groups

Catharine E. Fairbairn and Michael A. Sayette

University of Pittsburgh

Odd O. Aalen and Arnaldo Frigessi

University of Oslo

Catharine E. Fairbairn, Department of Psychology, University of Pittsburgh; Michael A. Sayette, Department of Psychology, University of Pittsburgh; Odd O. Aalen, Oslo Centre for Biostatistics and Epidemiology, Department of Biostatistics, University of Oslo; Arnaldo Frigessi, Oslo Centre for Biostatistics and Epidemiology, Department of Biostatistics, University of Oslo, and Research Support Services, Oslo University Hospital.

This research was supported by National Institutes of Health Grant (R01 AA015773) to Michael Sayette and National Science Foundation Graduate Research Fellowship (0753293-006) and National Science Foundation Graduate Research Opportunities Worldwide Grant (230165/F11) to Catharine Fairbairn. We thank John Dimoff, Michael Sayers, Bryan Heckman, Kasey Creswell, and the students and staff at the Alcohol and Smoking Research Laboratory for assistance collecting and coding data. We also thank Drs. Jeffrey Cohn, John Levine, Richard Moreland, Amanda Forest, and Thomas Kirchner for helpful comments regarding the group formation paradigm. Finally, we thank Dr. Harald Weedon-Fekjær and Dr. Terry Therneau for assistance with data analysis.

Correspondence concerning this article should be addressed either to Catharine Fairbairn, M.S., Department of Psychology, 3137 Sennott Square, University of Pittsburgh, 210 S. Bouquet St., Pittsburgh, PA 15260. Electronic mail: cef24@pitt.edu or to Michael Sayette, Ph.D.,

Department of Psychology, 3137 Sennott Square, University of Pittsburgh, 210 S. Bouquet St.,
Pittsburgh, PA 15260. Electronic mail: sayette@pitt.edu.

Abstract

Researchers have hypothesized that men gain greater reward from alcohol than women. However, alcohol-administration studies testing participants drinking alone have offered weak support for this hypothesis. Research suggests that social processes may be implicated in gender differences in drinking patterns. We examined the impact of gender and alcohol on “emotional contagion”—a social mechanism central to bonding and cohesion. Social drinkers (360 male, 360 female) consumed alcohol, placebo, or control beverages in groups of three. Social interactions were video recorded, and both Duchenne and non-Duchenne smiling were continuously coded using the *Facial Action Coding System*. Results revealed that Duchenne smiling (but not non-Duchenne smiling) contagion correlated with self-reported reward and typical drinking patterns. Importantly, Duchenne smiles were significantly less “infectious” among sober male versus female groups, and alcohol eliminated these gender differences in smiling contagion. Findings identify new directions for research exploring social-reward processes in the etiology of alcohol problems.

Keywords: alcohol, gender, emotional contagion, facial mimicry, social context

Individuals vary widely in their patterns of alcohol use, and researchers have long been interested in identifying individual difference criteria revealing those at risk to develop alcohol-related problems (Sher, Grekin, & Williams, 2005). In the search to identify predictors of alcohol abuse susceptibility, few individual-difference criteria have demonstrated as powerful a relationship with problematic drinking as has gender. Men show dramatically higher rates of alcohol-related problems than women. Men are 50% more likely than women to binge drink and twice as likely to report symptoms of alcohol abuse and dependence (SAMSA, 2012).

A longstanding hypothesis among alcohol researchers is that gender differences in susceptibility to alcohol use disorder may be traced to gender differences in alcohol reward sensitivity (Nolen-Hoeksema, 2004). According to this hypothesis, men would be expected to experience more subjective reinforcement from drinking alcohol than women (Sher, 1987; Wilson, 1988). Importantly, laboratory-based studies investigating gender differences in alcohol-related reinforcement have not produced consistent findings. While a handful of alcohol-administration studies report significant gender differences (Dougherty, Bjork, & Bennett, 1998; Mills & Bisgrove, 1983; Niaura, Nathan, Frankenstein, Shapiro, & Brick, 1987) many studies have not found significant differences across gender in response to alcohol consumption (e.g., Breslin, Mayward, & Baum, 1994; Josephs & Steele, 1990; Levenson, Oyama, & Meek, 1987; Mulvihill, Skilling, & Vogel-Sprott, 1997; Sayette, Breslin, Wilson, & Rosenblum, 1994; Sayette, Martin, Perrott, Wertz, & Hufford, 2001; Steele & Josephs, 1988). A variety of methodological factors may have contributed to these inconsistent findings, including lack of statistical power in many studies and a tendency to focus on self-report measures of mood.

One explanation for inconsistent effects of gender in the alcohol-administration literature is a failure to incorporate social context into experimental paradigms. Participants in laboratory-based alcohol studies tend to consume their study beverages in an isolated setting, with no other individual present in the room. These asocial drinking environments are likely poorly suited to an examination of differences according to gender. Based on decades of research examining patterns of gender differences across a variety of contexts, researchers have moved away from a static conceptualization of gender as a trait that predicts outcomes with equal regularity across setting (Wood & Eagly, 2010). Instead, gender is seen as a set of biological and societally determined predispositions that emerge selectively and dependent on elements of the proximal environment (Goffman, 1959). Importantly, researchers have observed only small and inconsistent gender differences along non-social outcomes (e.g., measures of verbal and quantitative ability), but dramatic differences between men and women when behavioral and emotional responses are observed in a social context (Eagly, 1995; Maccoby, 1990). Thus, theorists have identified “social interaction” as the context in which “the enactment of gender primarily takes place” (Deaux & Major, 1987, p. 370).

Several lines of alcohol research also suggest that social drinking paradigms could inform the understanding of gender differences in drinking patterns. Among both men and women, the overwhelming majority of drinking outside the laboratory occurs within the context of social interaction (Cahalan, Cisin, & Crossley, 1969; Demers et al., 2002; Single & Wortley, 1993; see Fairbairn & Sayette, in press). The formation and maintenance of social bonds has been identified as a fundamental human motive (Baumeister & Leary, 1995), and social factors exert considerable influence over emotions and behaviors. Across all drinking motives social enhancement motives show the most powerful and consistent differences according to gender,

with men drinking more to enhance social situations than women (e.g., Cooper, 1994; see Kuntsche, Knibbe, Gmel, & Engels, 2006) and studies examining social alcohol expectancies find similar gender effects (e.g., Rohsenow, 1983; see Wilson, 1988 for a review). All male drinking groups and all-male societies often show particularly high rates of hazardous drinking (Bartholow, Sher, & Krull, 2003; Bot, Engels, & Knibbe, 2005; Senchak, Leonard, & Greene, 1998; although see Rosenbluth, Nathan, & Lawson, 1978), and social factors may be more related to problematic drinking among men than among women (e.g., Mooney, Fromme, Kivlahan, & Marlatt, 1987). In sum, social environments are important in the study of problem drinking, and social factors may be implicated in gender differences in alcohol use disorders.

Gender, Social Bonding, and Emotional Contagion

A factor that differs dramatically across gender is the tendency to engage intimacy-building behaviors during social interactions. While research does not support the premise that men are less motivated than women to create close social relationships (Baumeister & Sommer, 1997), men display fewer affiliative behaviors during social interaction (Eagly & Wood, 1991). Socio-culturally defined gender roles are generally inconsistent with high levels of intimacy and affiliation displays among men, instead promoting displays of male competency and status (Briton & Hall, 1995; Deaux & LaFrance, 1998). Accordingly, research suggests that men are constrained in their displays of intimacy-building behaviors and demonstrate lower levels of social skill in affiliative social contexts. Men smile less during social interaction (LaFrance, Hecht, & Paluck, 2003), self-disclose less (Dindia & Allen, 1992) and demonstrate poorer emotion encoding (Hall, Coats, & LeBeau, 2005) and decoding abilities (Hall, 1978; McClure, 2000) than women. Women show greater responsiveness to the needs of others in close social relationships and offer more appropriate social support than men (Cultrona, 1996; Neff &

Karney, 2005). As men and women age from adolescence into adulthood, women benefit from larger and more supportive social networks (Antonucci & Akiyama, 1987). Thus, compared to women, men may not always feel free to engage social intimacy-building behaviors, and researchers have suggested that such constraints might carry negative consequences for their close social relationships (Baumeister & Sommer, 1997).

Emotional contagion¹ is an important mechanism through which feelings of affiliation and social connectedness are generated during social interaction (Chartrand & Lakin, 2013; Hatfield, Cacioppo, & Rapson, 1993; Hess & Fischer, 2013). In social contexts we unconsciously mimic the nonverbal expressions of our interaction partners and thereby “catch” their emotional states (Hatfield et al., 1995). Referred to by researchers as “social glue” (Lakin, Jefferis, Cheng, & Chartrand, 2003), emotional mimicry has been implicated in a variety of critical intimacy-building processes, including accurate decoding of emotional states of interaction partners (Niedenthal, 2007; Oberman, Winkielman, & Ramachandran, 2007; Stel & van Knippenberg, 2008). Individuals instructed to mimic the behaviors of their interaction partners report experiencing a more positive social experience (e.g., Stel & Vonk, 2010) and the relationship between emotional mimicry and social intimacy may be especially strong when smiling is the behavior being mimicked (Hess & Fischer, 2013).

While emotional contagion is common in social interaction, automatic mimicry responses may sometimes be inhibited (Chartrand & Lakin, 2013). For example, concerns with maintaining social status and perceived competence may constrain displays of mimicry (Kavanagh, Suhler, Churchland, & Winkielman, 2011; Lanzetta & Englis, 1989; Tiedens &

¹ “Emotional contagion” and “emotional mimicry” are closely related phenomena (Chartrand & Lakin, 2013)—the latter being considered the primary mechanism through which the former takes place (Hatfield et al., 1995). Both terms are used in this paper.

Fragale, 2003). Men are less likely to mimic emotional expressions than women, and research suggests that controlled inhibition of mimicry among men may account for these gender differences (Dimberg & Lundquist, 1990; Hatfield, Cacioppo, & Rapson, 1992; Sonny-Borgstrom, Jönsson, & Svensson, 2008). Experimental manipulations that reduce controlled inhibitive resources may sometimes facilitate the expression of emotional mimicry (van Leeuwen, van Baaren, Martin, Dijksterhuis, & Bekkering, 2009). A manipulation that has reliably impaired limited-capacity control processes is alcohol consumption (e.g., Fillmore, Vogel-Sprott, & Gavrilesco, 1999; Kirchner & Sayette, 2003; Steele & Josephs, 1990).

The Current Study

The current study is the first to our knowledge to test the impact of gender and alcohol consumption on emotional contagion. The dataset we use to examine this question has significant methodological advantages over prior studies of the effects of gender on alcohol response: 1) Participants in our study drink alcohol in a social context, a setting that is especially appropriate for an examination of gender differences (Deaux & Major, 1987); 2) Unlike many prior studies of gender differences in alcohol response, the current study includes both placebo and control comparison groups. In light of suggestions by researchers that men and women differ in their expectancies for alcohol-based reinforcement (Sher, 1987; Wilson, 1988), the ability to parse pharmacological from expectancy effects of alcohol represents a significant advantage²; and 3) During the social drinking period, facial muscle movements associated with smiling are continuously and unobtrusively assessed using the Facial Action Coding System (FACS) (Ekman, Friesen, & Hager, 2002). FACS is the most sophisticated and comprehensive

² The full 4-cell balanced placebo design has fallen out of favor in recent years, since participants provided with false information that they are not receiving alcohol are generally not deceived (Martin & Sayette, 1993; Testa et al., 2006). As a consequence, we employed a three cell design examining alcohol, placebo, and control conditions.

system for coding facial expression, serving as an objective and reliable measure of emotional state (Ekman & Rosenberg, 2005). For the present study, we focused on the most widely studied emotion-related expression in FACS, the Duchenne smile (e.g., Ekman, Davidson, & Friesen, 1990; Hess, Banse, & Kappas, 1995, Kirchner et al., 2006). Unlike other types of smiles, Duchenne smiles have been associated with “felt” rather than simply “displayed” emotion (Ekman & Rosenberg, 2005).

In order to assess emotional contagion in our group drinking setting we applied state-of-the-science statistical procedures capable of modeling dynamic social processes (Aalen, Borgan, & Gjessing, 2008). In an initial set of analyses, we examined the interaction of alcohol and gender by simply counting up the number of video frames in which all three members of a group smiled simultaneously (a group smile) during a social exchange (Sayette, Creswell, et al., 2012). Using this approach we found that, while both female gender and alcohol consumption were linked to greater levels of group smiling, alcohol did not interact with gender. These initial analyses provide a good starting place to examine alcohol’s social effects. Importantly, however, the static approach to behavioral assessment in Sayette et al. could not address emotional contagion, or the study of social coordination more generally. In particular: 1) The group level outcome in Sayette et al. (2012) did not permit examination of factors that vary within groups, such as gender. Accordingly, our initial analysis could not distinguish the effects of being a woman from that of being in a group with women and; 2) Our analytic approach did not model the sequence of individual-level events that preceded a group smile. Thus, group smiles that erupt almost instantaneously, sparked by any fleeting smile from an individual group member, are treated identically to group smiles that are drawn forth only after repeated and/or prolonged individual smiles.

In the current study we apply a statistical approach that, although rarely implemented within behavioral research, is nonetheless ideal for studying social interaction (Gardner & Griffin, 1989; Griffin & Gardner, 1989; Stoolmiller & Snyder, 2006). Here we model smiling using multilevel survival analysis, following the spread of smiles from one individual group member to the next. We focus on the simultaneous smile or mutual smile as a form of social coordination that carries important implications for social reward (Kirchner, Sayette, Cohn, Moreland, & Levine, 2006; Sayette, Creswell, et al., 2012). More specifically, we use survival analysis to examine the likelihood that a smile initiated by a single group member will 1) progress into a smile shared with another group member (a “mutual smile”) versus 2) end without evoking a responding smile (an “unreciprocated” smile). Unlike many commonly used behavioral analytic strategies (Bakeman & Gottman, 1997), survival analysis models not only the sequential ordering of behaviors but also their temporal duration and relative timing. Researchers have argued that a consideration of the relative timing of events is central to the study of emotional mimicry during unstructured interaction and also to the broader study of contagious processes (Diggle, 2006; Hove & Risen, 2009; Scheel, Aldrin, Frigessi, & Jansen, 2007). Our use of survival analyses allows us to ask a range of questions not possible using other approaches commonly applied to the study of social coordination including: 1) Are the effects of alcohol and gender on mutual smiling related to increases in the duration of initial smile or, instead, truly attributable to increases in social responsiveness?; 2) Are effects constant across the duration of the initial smile (i.e., do they arise immediately or only after a delay?); and 3) What is the effect of the smile initiator’s gender vs. the gender of the smile initiator’s group-mates? Within this paper, we examine these and other related issues, including the potential implications of “catching” a smile for social reward and typical drinking patterns.

We had two primary objectives in the present study. First, we aimed to use dynamic emotional assessment to examine the main effect of alcohol consumption on emotional contagion in a group drinking setting. Findings reported in our initial study (Sayette, Creswell, et al., 2012) suggested that alcohol increased incidence of simultaneous smiling, but left unanswered questions about potential mechanisms that might underlie these social effects (e.g., emotional contagion vs. individual-level processes). Second, we aimed to examine whether alcohol and gender interacted to predict the contagiousness of smiles. We hypothesized that gender differences in alcohol-related reinforcement would emerge. Specifically, we predicted that drinking alcohol would be associated with greater reward among men than among women, and that this reward would manifest selectively as a social, "catching" process. Accordingly, we predicted that, among sober groups, smiles would be less contagious in interactions containing more men than women, and that alcohol consumption would especially free men to engage in intimacy-building behaviors and "catch" the smiles of fellow group members. Thus, we predicted that alcohol would increase the contagiousness of smiles to a greater extent among men than among women in our social drinking setting.

Methods

Participants

As detailed elsewhere (Sayette, Creswell, et al., 2012), participants consisted of 720 healthy social drinkers aged 21-28 recruited via ads in local newspapers. Of these participants, 360 were male and 360 were female. Those who successfully completed an initial phone interview were invited to the Alcohol and Smoking Research Laboratory for a screening session. Following informed consent, exclusion criteria were assessed. Exclusion criteria included: medical conditions that contraindicated alcohol consumption, past alcohol abuse or dependence, as

indexed by DSM-IV, pregnancy in females, not being within 15% of ideal weight for height, and being uncomfortable with study drinking requirements. Eligible participants were invited to participate in the experiment (83% European-American, 11% African-American, 1% Hispanic, 2.5% Asian, and 2.5% other). Participants reported drinking 2-3 times/week and consuming 4.29 ($SD = 1.89$) drinks/occasion.

Procedure

Participants were informed that the purpose of the study was to measure alcohol's impact on cognitive performance. They were randomly assigned to groups of three, and groups were then assigned to an alcohol beverage (told alcohol, received alcohol), placebo beverage (told alcohol, received no alcohol), or control beverage (told no alcohol, received no alcohol) condition.

Within each of the three beverage conditions, groups were evenly distributed according to gender composition. Specifically, each beverage condition contained twenty all-female groups, twenty all-male groups, twenty groups with two females and one male, and twenty groups with two males and one female. Upon arriving in the lab, participants were casually and individually introduced to confirm that they were not previously acquainted (Kirchner et al., 2006).

[Additional participants were invited to each session to ensure that the three selected participants were unacquainted (see Sayette, Creswell, et al., 2012)]. Participants then provided a breath sample to assess blood alcohol content (BAC) and completed a variety of self-report mood and personality assessments.

The three participants were next seated at equidistant intervals around a round table. Cameras were positioned in all four corners of the room, and a microphone recorded conversation. Participants were originally told that the cameras were used to monitor their drink consumption and were later informed (see below) that the cameras recorded facial expressions.

Participants in the alcohol and placebo conditions were informed that they would be receiving alcohol and that the dose would be less than the legal driving limit. Drinks were mixed in front of all study groups (Rohsenow & Marlatt, 1981). The alcoholic beverage was 1 part 100 proof vodka and 3.5 parts Ocean Spray cranberry juice cocktail. In the placebo group, the glass was smeared with vodka, and a few drops of vodka were “floated” on the top of the beverage to increase credibility. We accounted for differences between men and women in rates of alcohol metabolism by adjusting the dose of alcohol according to gender. Males in the alcohol condition were administered a .82g/kg dose of alcohol, while females were administered a .74g/kg dose (Sayette et al., 2001). Participants remained seated for a total of 36-min while beverages were administered in three equal parts at 0-min, 12-min, and 24-min. They were instructed to drink their beverages evenly over the 12-min intervals and refrain from discussing how intoxicated they felt. Participants were otherwise not given instructions relevant to the social interaction—participants were ostensibly seated in the same room to facilitate drink administration and communication with the experimenter.

Immediately following drinking, participants’ BACs were recorded and they completed measures of mood and social bonding. Next they performed some additional cognitive tasks (see Sayette, Dimoff, Levine, Moreland, & Votruba-Drzal, 2012). After BAC was again assessed, placebo and control participants were debriefed, paid \$60, and allowed to leave. Participants in the alcohol condition remained until their BACs dropped below .025%. Before leaving, participants were informed that their behavior had been videotaped, and their consent to analyze the data was solicited (all participants agreed).

Participants’ facial expressions during the drinking period were later coded by FACS-certified personnel using Observer Video-Pro software (Noldus Information Technology, 2010).

The Observer system allows coders to time-stamp the start (onset) and stop (offset) of each facial muscle movement or action unit (AU) to preserve the flow and synchrony of the interaction. Each frame (1/30th of a second) of the interaction was manually evaluated by coders for the presence or absence of relevant facial AUs. Coders were blind to experimental condition. Video from each participant was independently coded so that the facial expressions of only one group member were visible to the coder at one time.

Measures

Typical Drinking Patterns: During their initial laboratory screening session, participants provided information concerning their typical drinking patterns. Specifically, they answered questions regarding the 1) Frequency of their alcohol use over the past 30 days (occasions/week) and 2) The average amount of alcohol (number of drinks) they consumed per drinking occasion.

Behavioral-Affective Display: We indexed “felt” emotional experience during the social interaction using the “Duchenne smile” (Ekman et al., 2002). Duchenne smiles include combined movement of the zygomaticus major (AU 12) and orbicularis oculi muscles (AU 6) (Ambadar, Cohn, & Reed, 2009; Ekman et al., 1990). “Displayed” emotion was indexed using the “non-Duchenne smile”—the movement of AU12 alone without the movement of AU6. Coders showed excellent inter-rater agreement for smiling (AU12, $\kappa = .84$; AU6, $\kappa = .88$).

Self-Reported Reward: Consistent with our past research (Fairbairn & Sayette, 2013) we indexed reward using self-report measures of mood and social bonding administered immediately after the interaction. We assessed social bonding using the Perceived Group Reinforcement Scale (PGRS). We assessed positive and negative mood using an 8-item Mood Measure (see Fairbairn & Sayette, 2013 for more details).

Data Analysis

Data Processing: Data were coded continuously throughout the 36 minute interaction with the exception of minutes 3-11 (only 29% of groups coded) and two additional minutes during which the experimenter entered the room to refill drinks. As before, we examine data from minutes 12-36 of the interaction—the period in which the effects of alcohol were hypothesized to be the strongest (Sayette, Creswell, et al., 2012). Approximately 34.9 million frames of behavioral data were coded. One group was excluded from analysis for technical reasons.

Beverage Condition: Beverage Condition was initially represented as a complete orthogonal set of contrast codes, the first (“Alcohol”) contrast comparing alcohol to both placebo and control conditions and the second (“placebo versus control”) contrast comparing placebo and control conditions (Cohen, Cohen, West, & Aiken, 2003). After confirming that there was empirical justification for collapsing across placebo and control conditions in these analyses (lack of significance of the placebo versus control contrast) we represent alcohol condition as a single dummy code.

Overview and Objectives: We aim to model the spread of smiles from one group member to the next, following smiles displayed by a single group member as they develop into smiles that are shared with other group members. To avoid redundancies within statistical models, we focus on smiles first initiated when no other group member is currently smiling. Given research highlighting the importance of timing in emotional contagion and mimicry processes, we use the word “infectiousness” in this paper in reference only to results of models that consider duration of the initial smile (i.e., survival analyses).

Many of the analyses implemented within this paper are well-established within science, but rarely applied within psychology (e.g., three-level nested frailty models). We therefore move

through our analyses in stages, beginning with the most simple and intuitive models and progressing through analytic approaches that increase in level of statistical complexity. As we arrive at each new stage in our analysis, we present a question that was left unanswered by previous (often more commonly implemented) statistical models together with a rationale for the next "stage" in our analysis. Specific information about data analytic strategies is presented immediately prior to the results produced by these models. Our analysis begins with simple event sequences at the level of the group, and ends with models that examine within-group predictors and incorporate a complex consideration of event duration.

Results

Beverage Manipulation Check and Baseline Comparisons

BACs and measures of subjective intoxication appear in Table 1. Participants drinking alcohol were on the rising limb of the BAC curve with a BAC of just above .06% immediately following the interaction period. All placebo and alcohol participants estimated that they had consumed at least 1 oz. of vodka. Consistent with our prior studies (e.g., Sayette et al., 2001), placebo participants reported experiencing some level of intoxication, more than control participants and less than alcohol participants. Neither gender nor typical drinking patterns significantly affected BACs or subjective levels of intoxication. As reported elsewhere, there were no differences in baseline personality or mood variables across drink conditions (see Sayette, Creswell, et al., 2012).

Event-Level Analysis

We begin by exploring the spreading of smiles within groups at the level of the "event." In other words, we test the probability of a single smile developing into a mutual smile (versus an "unreciprocated smile") without considering the *duration* of smiles under examination.

Consistent with the notion that alcohol enhanced social bonding, event probabilities displayed in Table 2 suggest that smiles initiated in groups consuming alcohol were more likely to lead to mutual smiles (49.6%) than were those initiated in the placebo (45.6%) or control (44.6%) conditions, with the latter two conditions not differing. Further, irrespective of drink condition, smiles initiated in groups consisting of all males were less likely to develop into mutual smiles (42.6%) than were those initiated in all-female groups (48.6%). Finally, a glance at these descriptive statistics suggests that the effect of alcohol on increasing transitions to mutual smiles may be larger in groups containing more males. Hierarchical logistic regression models (Kenny, Kashy, & Cook, 2006; Raudenbush & Bryk, 2002) confirm many of these observations. Alcohol significantly increased the likelihood that a smile would develop into a mutual smile, $B = 0.219$, $OR = 1.245$, $t = 3.260$, $p = 0.001$, as did the number of women included in the group, $B = 0.092$, $OR = 1.096$, $t = 3.160$, $p = 0.002$. Further, there was a marginally significant alcohol by group gender composition interaction, $B = -0.116$, $OR = 0.890$, $t = -1.920$, $p = 0.055$, suggesting that this main effect of alcohol was driven by its effects on men. The distinction between placebo and control conditions did not emerge as significant in any analysis, p 's $> .8$.

Survival Analysis

The event-level analysis described above provides a useful starting point in our examination of social coordination. Unlike analyses implemented in our previous work (Sayette, Creswell, et al., 2012), this event-level approach considers the sequencing of smiles within groups, accounting not only for the number of mutual smiles displayed during the social interaction, but also the number of individual smiles that were left “unreciprocated.” Analyses that examine social interactions as sequences of events represent the single most popular framework for the examination of social exchange (e.g., sequential analysis, Bakeman &

Gottman, 1997). Importantly, however, this event-level approach ignores a substantial amount of valuable information contained in the data. An event-level analysis does not consider the duration of the behavior under examination. A smile that develops into a mutual smile after 0.2 seconds, for example, is treated identically to a smile that develops into a mutual smile after 8 seconds (Griffin & Gardner, 1989). In the current study, as is increasingly common in studies of social interaction, data were not collected at the level of the event, but instead behavior was continuously coded in time, producing a dataset so precise as to carry information down to the level of the frame (1/30th of a second). Effectively “throwing out” event duration information might in some cases undermine power to detect significant effects (Green & Symons, 1983). Further, a behavior displayed over a prolonged period may have different effects than this same behavior displayed only fleetingly (Ekman, 2009), and therefore, depending on the theoretical focus of the study, a strictly event-level analysis may lead to misconceptions concerning the nature of the social process under examination.

Given the focus on contagious processes in the current project, concluding our analysis at the level of the event would seem to pose problems. In any examination seeking to determine the level of “infectiousness” of a given agent, information concerning intensity of exposure to the proposed pathogen is highly relevant (Diggle, Heagerty, Liang, & Zeger, 1994; Diggle, 2006; Scheel et al., 2007). Importantly, in the current study, smiles in the alcohol condition lasted on average one second longer (4.5sec) than did smiles in the no alcohol conditions (3.47sec placebo, 3.48sec control), $B = 1.048$, $t = 6.99$, $p < .0001$. Thus, any main effects or interactions involving alcohol could potentially have emerged simply as a result of these increased smile durations. In other words, we might observe increases in mutual smiling with alcohol not because alcohol

increases the innate “infectiousness” of smiles per unit time, but simply because exposure to the initial smile was more prolonged.

In contrast to event-focused approaches, survival analysis considers not only the *sequence* of events but also event *duration* (Gardner & Griffin, 1989; Griffin & Gardner, 1989; Stoolmiller & Snyder, 2006). The quantity that is studied within a survival framework is the “hazard” or the probability of a given outcome occurring per unit time under examination. In the results reported below, the hazard is represented by the abbreviation $Exp(B)$, and can be interpreted as a form of “relative risk” across levels of the predictors. At this stage of our analysis, we employed a “competing risks” frailty survival model accounting for clustering of recurrent events at the level of the group (Allison, 2012).

Results of Survival Models: Group gender composition significantly impacted the hazard of a smile developing into a mutual smile, $B = 0.088$, $Exp(B) = 1.092$, $SE(B) = .022$, $p < .0001$. Introducing each woman into a group resulted in a 9.2% increase in transitions into mutual smiles per unit time that the initial smile was displayed (see Figure 1). In contrast to the event-level analysis presented above, however, the survival model produced no significant main effect of alcohol, $p = .712$. Accounting for each unit time that a smile was displayed, there was no main effect of alcohol on the probability that a smile would develop into a mutual smile. When considered together with prior analyses, these results suggest that broad alcohol-related increases in mutual smiling are attributable to alcohol’s tendency to prolong the duration of individual (initial) smiles. Survival analysis reveals that, per unit time that the initial smile was displayed, alcohol had no impact on the likelihood that the initial smile would be reciprocated.

Importantly, the effect of alcohol on mutual smile hazard varied depending on group gender composition. Alcohol significantly increased the contagiousness of smiles selectively

among-all male groups. Survival models indicate a significant alcohol by group gender composition interaction, $B = -0.109$, $Exp(B) = 0.897$, $SE(B) = .045$, $p = .015$. Alcohol did not significantly increase the hazard of transition to mutual smiling when groups contained any women. In contrast, we found that alcohol was associated with an increase in hazard of transition to mutual smiling among groups consisting of all males. When groups consisted of only men, alcohol was associated with an increase of 21% in transitions to mutual smiling per unit time that the initial smile was displayed, $B = 0.192$, $Exp(B) = 1.211$, $SE(B) = .086$, $p = .025$. Thus, selectively among all-male groups, alcohol-related increases in mutual smiling were not solely attributable to increased initial smile duration. Sober all-male groups showed significantly fewer mutual-smile transitions when compared to sober groups containing women, $B = -0.128$, $Exp(B) = 0.879$, $SE(B) = .027$, $p < .0001$, and alcohol eliminated this male deficit in mutual smiling hazard. Again, we found no significant distinction between placebo and control conditions in any of the above analyses, p 's $> .42$.

Is this Really Contagion?

Thus far, our analyses have considered only situations in which one group member has already initiated a smile. We have not considered situations in which no group members are currently smiling. If the gender by alcohol interaction identified above manifests as a result of a purely “contagious” process, passed from one group member to the next, we would expect this interaction to emerge selectively in situations when one group member has initiated a smile. Results reported to this point do not rule out the possibility that the outcome we refer to as smile “contagiousness” simply reflects predilection to smile, regardless of whether another group member is already smiling.

To address this question, we examined whether the alcohol by gender interaction emerged to predict smiling hazard in situations in which no other group members are currently smiling ("non-infection" model). Results derived from this "non-infection" model can then be compared with results produced by the "infection models" reported above, which consider situations in which another group member is already smiling. In this case, we constructed two separate models in order to allow the shape of the hazard to vary across these distinct situations. Importantly, within the "non-infection" framework the alcohol by group gender composition interaction does not approach significance, $B = 0.002$, $Exp(B) = 1.002$, $SE(B) = 0.047$, $p = 0.960$. Thus, it appears that alcohol's tendency to increase mutual smiling selectively in all-male groups is not explained by overall increases in smiling, and instead appears more similar to a "contagious" process that manifests upon exposure to a smile from another group member.

Do Results Vary across Smile Duration?

A unique aspect of survival analysis is its capability for addressing sophisticated hypotheses concerning the effect of variations in event timing on hazards (Allison, 2012). Within psychology, information concerning whether an effect emerges instantaneously versus after a delay can explicate the mechanisms underlying this process (e.g., automatic versus controlled) (Greenwald, McGhee, & Schwartz, 1998). Here, we explored whether effects of gender and alcohol on transitions to mutual smiling varied as a function of the time *after* onset of the initial smile. We uncovered a significant group gender by alcohol by smile duration interaction, $B = -.041$, $Exp(B) = .960$, $SE(B) = .018$, $p = .023$. Among groups not consuming alcohol, there was a significant smile duration by group gender composition interaction, $B = .027$, $Exp(B) = 1.028$, $SE(B) = .012$, $p = .019$. Sober groups showed similar rates of transition to mutual smiles in an *instantaneous or immediate* fashion regardless of gender composition, but

after a *delay*, sober all-male groups showed lower hazards to mutual smile than groups containing more females. Alcohol acted to eliminate this deficit for males in non-instantaneous (delayed) mutual smile hazard, $p = .986$.

Moving Beyond the Group

To this point, our analyses have been conducted strictly at the level of the group. However, unlike analyses implemented in many previous studies of behavioral coordination (e.g., Sayette, Creswell, et al., 2012), survival analysis allows for the differentiation of the individuals acting within each group. Specifically, we can disentangle the effects of the gender of the individual who initiated the smile from the gender of this individual's group-mates. Thus, using Kenny and colleagues' actor-partner interdependence model (Kenny & Garcia, 2012), we sought to differentiate the effects of being a man from the effects of being in a group with men on hazard to mutual smiling.

Applications of survival analysis to social interaction have not previously moved beyond the level of the group, accounting for nesting at only two levels of analysis—within-group and between groups (Stoolmiller & Snyder, 2006). However, the three-level (“nested”) frailty model has been well-studied in fields outside of psychology, and software is now readily available for implementing such analyses (Sastry, 1997; Therneau, 2011). We conducted a “nested” frailty model within the R software program *coxme* (Therneau, 2011). Results indicate that the gender of the individual who initiates the smile does not impact the hazard that a smile will transition into a mutual smile. In other words, this study did not find evidence that smiles displayed by women are more “infectious,” and we found no moderating influence of alcohol on this effect, p 's $> .52$. *Instead, the group-level gender composition effects outlined above appear to be driven by the gender of the smile initiator's group-mates (potential smile responders).* Hazard to

mutual smile was 23% lower if smiles were initiated in groups with two male group-mates versus groups with one or two female group-mates, $B = -0.257$, $Exp(B) = .773$, $SE(B) = .048$, $p < .0001$. If the smiler's group-mates included at least one female, then hazard to mutual smile reached equivalent levels to those observed in groups including two female group-mates, $p = .18$.

Finally, there was a highly significant interaction between group-mates' gender and alcohol condition, $B = 0.373$, $Exp(B) = 1.452$, $SE(B) = 0.098$, $p = .0001$. When both group-mates were men, alcohol increased the hazard of a smile developing into a mutual smile by 30%, $B = 0.263$, $Exp(B) = 1.302$, $SE(B) = 0.083$, $p = .002$ (Figure 2). In contrast, there was no significant effect of alcohol on hazard to mutual smiles when at least one group-mate was a female. These results suggest that alcohol selectively boosts the chances that men will "catch" a smile, and may have comparatively little effect on smile infection passed to women.

Clinical and Reward Implications (What Does it Mean to "Catch" a Smile?)

We next examined the potential implications of smile infection for alcohol-related reward and reinforcement value. First, we explored the relationship between smile infectiousness and reward by examining associations between mutual smile hazard and self-reported indexes of mood and social bonding administered after the interaction. Similar to gender findings reported above, effects emerged primarily in models considering attributes of the smile initiator's group-mates. Specifically, increased probability of smile "catching" corresponded to significant increases in positive mood, $B = 0.065$, $Exp(B) = 1.067$, $SE(B) = 0.019$, $p = .0005$, and social bonding, $B = 0.023$, $Exp(B) = 1.023$, $SE(B) = 0.011$, $p = 0.046$, together with decreases in negative mood, $B = -0.095$, $Exp(B) = 0.909$, $SE(B) = 0.027$, $p = .0004$. Some significant effects and trends also emerged with respect to the smile initiator's mood, with increased mutual smile hazard predicting significantly decreased negative mood, $B = -0.076$, $Exp(B) = 0.927$, $SE(B) =$

0.038, $p = .0045$, and tending to increase positive mood, $B = 0.044$, $Exp(B) = 1.045$, $SE(B) = 0.025$, $p = 0.083$, although no significant effect emerged with respect to social bonding, $B = 0.022$, $Exp(B) = 1.022$, $SE(B) = 0.016$, $p = 0.180$.

Since propensity to “catch” a smile appears to correspond to reward, sensitivity to alcohol’s impact on smile infection might reinforce drinking and thus might be used to predict individual differences in typical drinking patterns. In line with this supposition, analyses revealed a significant interaction between alcohol condition and group-mates’ typical drinking patterns, $B = 0.091$, $Exp(B) = 1.096$, $SE(B) = 0.033$, $p = 0.006$. Among groups consuming alcohol, a smile was significantly more likely to be “caught” if the smile initiator’s group-mates were heavier vs. lighter drinkers. More specifically, within the alcohol condition, there was a significant positive association between mutual smile hazard and group-mates’ typical drinking quantity, $B = 0.055$, $Exp(B) = 1.056$, $SE(B) = 0.026$, $p = 0.037$, such that a 5.6% increase in hazard of mutual smile corresponded to an increase of 1 standard drink per typical drinking occasion. In contrast, drinking patterns did not significantly predict mutual smile hazard in the placebo and control conditions. The typical drinking patterns of the smile initiator him/herself did not significantly predict hazard of mutual smile in any of the beverage conditions, $p = .72$. The effect of group-mates’ typical drinking patterns was most pronounced in models examining drinking quantity (average drinks/occasion, results reported above), while the effect of drink frequency alone did not reach significance, $p = .69$. Of note, models examining total alcohol consumed (quantityXfrequency) did reveal a significant alcohol condition by group-mates’ drinking pattern interaction, $B = 0.017$, $Exp(B) = 1.017$, $SE(B) = 0.007$, $p = 0.016$. While gender was a powerful predictor of drinking quantity among participants in our study, $t = 9.500$, $p < .0001$, with men consuming significantly more alcohol per drinking occasion than women, the effect of typical

drinking patterns emerged as significant even in models controlling for all moderated and unmoderated effects of gender. Interestingly, associations of smile hazard with group-mates' drinking patterns appeared to be specific to the "infection" model (when one group member is already smiling), and did not emerge as significant within the "non-infection" framework discussed earlier, $p = .98$. Thus, results suggest that the "infectiousness" of smiles in a group drinking setting could have specific and clinically meaningful implications for social reward and alcohol-related reinforcement.

What about non-Duchenne Smiles?

To test whether effects examined here were specific to "emotion"-related expressions—if effects could truly be called "emotional" contagion—we examined whether significant findings reported above would generalize to models examining non-Duchenne smiles. Results suggested that "catching" a non-Duchenne smile was not associated with self-reported reward, p 's $> .28$, and did not interact with alcohol condition to predict typical drinking patterns, $p = .19$. Further, unlike Duchenne smiles, gender effects did not vary significantly by alcohol condition in non-Duchenne smiling infection models, $p = .11$.

Discussion

For years the proposition that alcohol consumption differentially affects emotion for men and women has been debated. A typical conclusion is that gender ought to influence the effects of alcohol, but the experimental findings are stubbornly complex if not downright inconsistent (Sher, 1987; Sutker, Allain, Brantley, & Randall, 1982). We have argued here that a vigorous investigation of the moderating impact of gender on the alcohol-emotion relation requires consideration of social context. While at first glance this position seems to be fairly straightforward, in fact it is at odds with the majority of experimental alcohol research.

Indeed, emotions and motivations associated with social drinking are not widely believed to carry serious implications for the etiology of alcohol use disorder. Unlike individuals consuming alcohol while alone, those who drink to intoxication in a social setting do so with the implied social sanction and, perhaps, facilitation of those individuals in their proximal environment. Thus, while the affective experience of solitary drinkers is presumed to be truly “intrinsic” and “fundamental,” we assume that the emotional and motivational forces that drive alcohol consumption in a social setting are “watered-down” and “secondary” in comparison. Consistent with such assumptions, researchers seldom incorporate social drinking paradigms into alcohol-administration studies, and drinking in social settings has rarely been subject to serious scientific scrutiny. These assumptions and practices not only overlook the fact that, regardless of an individual’s problem drinking status, the majority of drinking occurs in social settings but also disregard research suggesting that social motives are among the most “fundamental” that humans possess. Indeed, our own experience dictates that emotions that manifest during social interactions can be overwhelming. Certainly the reward we experience when we share a genuine smile of enjoyment with another individual can represent a powerful motivational force. It is in this context that the present study, which integrates research and methods across several domains of psychology, should be considered.

Using dynamic measures of affective experience and a multilevel statistical approach, we examined the impact of alcohol consumption and group gender composition on the spreading of smiles within groups. Specifically, we tested the effect of alcohol and group gender on the probability that an initial smile would develop into a mutual smile. Our initial event-level set of analyses focused on the likelihood that an initial smile, *irrespective of its duration*, would develop into a mutual smile. This analysis represents the dominant statistical approach to the

examination of social interaction (Bakeman & Gottman, 1997). Results indicated that groups consuming alcohol were more likely to reciprocate a smile than were groups consuming nonalcoholic beverages. That is, alcohol increased the probability that an initial smile would develop into a mutual smile. Further, this effect of alcohol appeared to be more pronounced for men than women.

By themselves, these initial analyses leave unresolved critical questions regarding the nature of this gender by alcohol interaction. Indeed, had we stopped at this stage in our analysis we might have mistakenly assumed that alcohol consumption enhanced social responsiveness and emotional contagion, with alcohol qualitatively altering the infectiousness of initial smiles. Alternatively, a mutual smile may occur simply because the initial smile lingers in an unreciprocated state for a long enough time that eventually another group member responds. The increased likelihood of a mutual smile would not be due to any fundamental change in the inherent infectiousness of the initial smile. The event-level analysis cannot disentangle these conceptually distinct possibilities.

Survival analysis allows for a complex consideration of not only event sequence but also event duration. When we apply survival analysis to our data, we find that the main effect of alcohol on increasing mutual smiling is fully accounted for by increases in initial smile duration. These results suggest that alcohol's global tendencies to increase mutual smiling in social settings may not result from increases in emotional contagion or social responsiveness, but instead may stem from individual-level effects on something akin to "social bravery" (Fairbairn, Sayette, Levine, Cohn, & Creswell, 2013; Fairbairn & Sayette, in press). Alcohol appears to increase our willingness to allow our "true smiles" to linger longer in a social setting, and, as a result, increases the likelihood that they will ultimately meet with a responding smile.

When considering the impact of gender, our survival analyses indicate that men derive more reward from alcohol than women, and that this reward manifested as a specifically social, "catching" process. Among men, alcohol-related increases in mutual smiling are not entirely accounted for by increases in the duration of the initial smile. Alcohol increased smiling "infectiousness" selectively among all-male groups but not among groups containing females. Sober all-male groups displayed lower rates of smile contagiousness than groups containing females, and alcohol consumption eliminated this male-group deficit in smiling infection. Further analyses reveal that this significant group-level interaction was not driven by the gender of the smile initiator but was instead driven by the gender of the smile initiator's group-mates--the gender of those who might be "infected" by a smile. Thus, results suggest that alcohol selectively enhanced the probability that a man will "catch" a smile in a social drinking context.

In addition, we used survival analysis to examine whether effects of predictors on the probability of mutual smiling varied across the duration of the initial smile. We found that sober male deficits in hazard to mutual smiling did not arise instantaneously, but instead became more pronounced as initial smile duration progressed. Prior research has found that men and women do not differ in levels of emotional mimicry when viewing pictures of emotional faces at subliminal exposure times, but instead that gender differences emerge selectively at superliminal levels of exposure (Sonny-Borgstrom et al., 2008). Consequently, when considered in light of prior research, our results suggest a role for more deliberative processes in gender differences in emotional mimicry. Consistent with research documenting alcohol's tendency to reduce controlled inhibition, results of this study hint that alcohol may reduce gender differences in emotional contagion by disrupting non-automatic processes that otherwise would constrain mimicry among sober men.

Finally, survival analyses revealed links between smile infectiousness and indexes of reward and alcohol-related reinforcement. Increased probability of “catching” a smile was associated with enhanced reported mood and perceived social outcomes. Further, selectively in the alcohol condition, individuals who were more likely to be “infected” by a smile were also more likely to report being heavier drinkers outside the laboratory. The relationship between drinking pattern and smile infection was independent of gender and, when considered together with self-reported reward findings, suggests that smile infection could represent an important indicator of alcohol-related reinforcement and a mechanism supporting drinking

Limitations of this study should be noted. First, we employed a single moderate dose of alcohol and tested the responses of individuals while on the ascending limb of the BAC curve. Future studies should test the generalizability of these results to higher and lower doses of alcohol and to individuals whose BACs are descending. Second, the current research examined emotional contagion by using a measure of positive emotion (the smile) and a non-threatening social interaction paradigm. Further research might explore alcohol’s impact on the contagion of negative emotion during threatening or stressful social interaction. Third, we examined alcohol’s impact on smiling only in a social setting and not among participants consuming alcohol while alone. While previous research does not support the notion that alcohol increases smiling duration among participants drinking in isolation (Ruch, 1994; Sayette et al., 2001), we are nonetheless unable to definitively claim that effects observed in this paper are uniquely social phenomena or, for example, that alcohol’s effects on increasing smile duration represent a manifestation of “social bravery.” Further, here we examine effects of alcohol among groups of individuals who were not previously acquainted, and future research should examine the generalizability of these findings to groups at all stages of social integration (Fairbairn &

Sayette, in press). Fourth, here we identify only one potential source of alcohol-related reinforcement. We do not claim that smile infectiousness is the only or even the most important manifestation of the reward that might be derived from drinking alcohol. Other factors—e.g., the duration of the mutual smile—may also support alcohol-related reward and reinforcement-value. Finally, the present paradigm mirrors common real-world situations in which all members of a social group consume alcohol together. Future research would be indicated to extend these findings by varying drink condition within groups.

In summary, the present study integrated theory and methods from diverse fields to examine the impact of alcohol on social experience. Results provide evidence suggesting that the social rewards associated with alcohol consumption are especially pronounced for male drinkers. Findings indicate a novel yet fundamental source of reward for male social drinkers, and provide new directions for research exploring the role of social factors in the etiology of alcohol use disorder.

Authorship

C.E.F developed the study concept. M.A.S. developed the study design and participated in data collection. C.E.F. performed the data analysis and interpretation under the supervision of A.F. and O.O.A. C.E.F. drafted the paper, and M.A.S. provided critical revisions. All authors approved the final version of the paper for submission.

References

- Aalen, O. O., Borgan, O., & Gjessing, H. (2008). *Survival and event history analysis: A process point of view*. New York: Springer.
- Allison, P. D. (2012). *Survival analysis using SAS: A practical guide*. Cary, NC: SAS Institute.
- Ambadar, Z., Cohn, J. F., & Reed, L. I. (2009). All smiles are not created equal: Morphology and timing of smiles perceived as amused, polite, and embarrassed/nervous. *Journal of Nonverbal Behavior*, *33*, 17–34. doi:10.1007/s10919-008-0059-5
- Antonucci, T. C., & Akiyama, H. (1987). An examination of sex differences in social support among older men and women. *Sex Roles*, *17*, 737–749. doi:10.1007/BF00287685
- Bakeman, R., & Gottman, J. M. (1997). *Observing interaction: An introduction to sequential analysis* (2nd ed.). New York: Cambridge University Press.
- Bartholow, B. D., Sher, K. J., & Krull, J. L. (2003). Changes in heavy drinking over the third decade of life as a function of collegiate fraternity and sorority involvement: A prospective, multilevel analysis. *Health Psychology*, *22*, 616–626. doi:10.1037/0278-6133.22.6.616
- Baumeister, R. F., & Leary, M. R. (1995). The need to belong: Desire for interpersonal attachments as a fundamental human motivation. *Psychological Bulletin*, *117*, 497–529. doi:10.1037/0033-2909.117.3.497
- Baumeister, R. F., & Sommer, K. L. (1997). What do men want? Gender differences and two spheres of belongingness: Comment on Cross and Madson (1997). *Psychological Bulletin*, *122*, 38–44. doi:10.1037/0033-2909.122.1.38

- Bot, S. M., Engels, R. C. M. E., & Knibbe, R. A. (2005). The effects of alcohol expectancies on drinking behaviour in peer groups: Observations in a naturalistic setting. *Addiction, 100*, 1270–1279. doi:10.1111/j.1360-0443.2005.01152.x
- Breslin, F. C., Mayward, M., & Baum, A. (1994). Effect of stress on perceived intoxication and the blood alcohol curve in men and women. *Health Psychology, 13*, 479–487. doi:10.1037/0278-6133.13.6.479
- Briton, N. J., & Hall, J. A. (1995). Beliefs about female and male nonverbal communication. *Sex Roles, 32*, 79–90. doi:10.1007/BF01544758
- Cahalan, D., Cisin, I. H., & Crossley, H. M. (1969). *American drinking practices: A national study of drinking behavior and attitudes*. New Brunswick, NJ: Rutgers Center of Alcohol Studies.
- Chartrand, T. L., & Lakin, J. L. (2013). The antecedents and consequences of human behavioral mimicry. *Annual Review of Psychology, 64*, 285–308. doi:10.1146/annurev-psych-113011-143754
- Cohen, J., Cohen, P., West, S. G., & Aiken, L. S. (2003). *Applied multiple regression/correlation analysis for the behavioral sciences* (3rd ed.). Hillsdale, NJ: Routledge Academic.
- Cooper, M. L. (1994). Motivations for alcohol use among adolescents: Development and validation of a four-factor model. *Psychological Assessment, 6*, 117–128. doi:10.1037/1040-3590.6.2.117
- Cultrona, C. E. (1996). *Social support in couples*. Thousand Oaks, CA: Sage.
- Deaux, K., & LaFrance, M. (1998). Gender. In D. Gilbert, S. T. Fiske, & G. Lindzey (Eds.), *Handbook of social psychology* (Vol. 4). New York: Random House.

- Deaux, K., & Major, B. (1987). Putting gender into context: An interactive model of gender-related behavior. *Psychological Review*, *94*, 369–389. doi:10.1037/0033-295X.94.3.369
- Demers, A., Kairouz, S., Adlaf, E., Gliksman, L., Newton-Taylor, B., & Marchand, A. (2002). Multilevel analysis of situational drinking among Canadian undergraduates. *Social Science & Medicine*, *55*, 415–424. doi:10.1016/S0277-9536(01)00258-1
- Diggle, P. J. (2006). Spatio-temporal point processes, partial likelihood, foot and mouth disease. *Statistical Methods in Medical Research*, *15*, 325–336. doi:10.1191/0962280206sm454oa
- Diggle, P. J., Heagerty, P., Liang, K., & Zeger, S. (1994). *Analysis of longitudinal data*. Oxford: Clarendon Press.
- Dimberg, U., & Lundquist, L. (1990). Gender differences in facial reactions to facial expressions. *Biological Psychology*, *30*, 151–159. doi:10.1016/0301-0511(90)90024-Q
- Dindia, K., & Allen, M. (1992). Sex differences in self-disclosure: A meta-analysis. *Psychological Bulletin*, *112*, 106–124. doi:10.1037/0033-2909.112.1.106
- Dougherty, D. M., Bjork, J. M., & Bennett, R. H. (1998). Effects of alcohol on rotary pursuit performance: A gender comparison. *The Psychological Record*, *48*, 393–405.
- Eagly, A. H. (1995). The science and politics of comparing women and men. *American Psychologist*, *50*, 513–520. doi:10.1037/0003-066X.45.4.513
- Eagly, A. H., & Wood, W. (1991). Explaining sex differences in social behavior: A meta-analytic perspective. *Personality and Social Psychology Bulletin*, *17*, 306–315. doi:10.1177/0146167291173011
- Ekman, P. (2009). *Telling Lies: Clues to Deceit in the Marketplace, Politics, and Marriage, Third Edition (Revised Edition.)*. W. W. Norton & Company.

- Ekman, P., Davidson, R. J., & Friesen, W. V. (1990). The Duchenne smile: Emotional expression and brain physiology: II. *Journal of Personality and Social Psychology*, *58*, 342–353. doi:10.1037/0022-3514.58.2.342
- Ekman, P., Friesen, W. V., & Hager, J. C. (2002). *Facial Action Coding System*. Salt Lake City, UT: Network Information Research.
- Ekman, P., & Rosenberg, E. L. (2005). *What the face reveals: Basic and applied studies of spontaneous expression using the Facial Action Coding System (FACS)* (2nd ed.). New York, NY: Oxford University Press.
- Fairbairn, C. E., & Sayette, M. A. (in press). A social-attributational analysis of alcohol response. *Psychological Bulletin*.
- Fairbairn, C. E., & Sayette, M. A. (2013). The effect of alcohol on emotional inertia: A test of alcohol myopia. *Journal of Abnormal Psychology*, *122*, 770–781. doi:10.1037/a0032980
- Fairbairn, C. E., Sayette, M. A., Levine, J. M., Cohn, J. F., & Creswell, K. G. (2013). The effect of alcohol on the emotional displays of whites in interracial groups. *Emotion*, *13*, 468–477. doi:10.1037/a0030934
- Fillmore, M. T., Vogel-Sprott, M., & Gavrilesco, D. (1999). Alcohol effects on intentional behavior: Dissociating controlled and automatic influences. *Experimental and Clinical Psychopharmacology*, *7*, 372–378. doi:10.1037/1064-1297.7.4.372
- Gardner, W., & Griffin, W. A. (1989). Methods for the analysis of parallel streams of continuously recorded social behaviors. *Psychological Bulletin*, *105*, 446–455. doi:10.1037/0033-2909.105.3.446
- Goffman, E. (1959). *The presentation of self in everyday life*. Garden City, NY: Double Day Anchor Books.

- Green, M. S., & Symons, M. J. (1983). A comparison of the logistic risk function and the proportional hazards model in prospective epidemiologic studies. *Journal of Chronic Diseases, 36*, 715–723. doi:10.1016/0021-9681(83)90165-0
- Greenwald, A. G., McGhee, D. E., & Schwartz, J. L. (1998). Measuring individual differences in implicit cognition: The implicit association test. *Journal of Personality and Social Psychology, 74*, 1464–1480. doi:10.1037/0022-3514.74.6.1464
- Griffin, W. A., & Gardner, W. (1989). Analysis of behavioral durations in observational studies of social interaction. *Psychological Bulletin, 106*, 497–502. doi:10.1037/0033-2909.106.3.497
- Hall, J. A. (1978). Gender effects in decoding nonverbal cues. *Psychological Bulletin, 85*, 845–857. doi:10.1037/0033-2909.85.4.845
- Hall, J. A., Coats, E. J., & LeBeau, L. S. (2005). Nonverbal behavior and the vertical dimension of social relations: A meta-analysis. *Psychological Bulletin, 131*, 898–924. doi:10.1037/0033-2909.131.6.898
- Hatfield, E., Cacioppo, J. T., & Rapson, R. L. (1992). Primitive emotional contagion. *Review of Personality and Social Psychology, 14*, 151–177.
- Hatfield, E., Cacioppo, J. T., & Rapson, R. L. (1993). Emotional contagion. *Current Directions in Psychological Science, 2*, 96–99. doi:10.1111/1467-8721.ep10770953
- Hess, U., & Fischer, A. (2013). Emotional mimicry as social regulation. *Personality and Social Psychology Review, 17*, 142–157. doi:10.1177/1088868312472607
- Hove, M. J., & Risen, J. L. (2009). It's all in the timing: Interpersonal synchrony increases affiliation. *Social Cognition, 27*, 949–960. doi:10.1521/soco.2009.27.6.949

- Josephs, R. A., & Steele, C. M. (1990). The two faces of alcohol myopia: Attentional mediation of psychological stress. *Journal of Abnormal Psychology, 99*, 115–126.
doi:10.1037/0021-843X.99.2.115
- Kavanagh, L. C., Suhler, C. L., Churchland, P. S., & Winkielman, P. (2011). When It's an Error to Mirror: The Surprising Reputational Costs of Mimicry. *Psychological Science, 22*, 1274–1276. doi:10.1177/0956797611418678
- Kenny, D. A., & Garcia, R. L. (2012). Using the actor–partner interdependence model to study the effects of group composition. *Small Group Research, 43*, 468–496.
doi:10.1177/1046496412441626
- Kenny, D. A., Kashy, D. A., & Cook, W. L. (2006). *Dyadic data analysis*. New York, NY: The Guilford Press.
- Kirchner, T. R., & Sayette, M. A. (2003). Effects of alcohol on controlled and automatic memory processes. *Experimental and Clinical Psychopharmacology, 11*, 167–175.
- Kirchner, T. R., Sayette, M. A., Cohn, J. F., Moreland, R. L., & Levine, J. M. (2006). Effects of alcohol on group formation among male social drinkers. *Journal of Studies on Alcohol, 67*, 785–793.
- Kuntsche, E., Knibbe, R., Gmel, G., & Engels, R. (2006). Who drinks and why? A review of socio-demographic, personality, and contextual issues behind the drinking motives in young people. *Addictive Behaviors, 31*, 1844–1857. doi:10.1016/j.addbeh.2005.12.028
- LaFrance, M., Hecht, M. A., & Paluck, E. L. (2003). The contingent smile: A meta-analysis of sex differences in smiling. *Psychological Bulletin, 129*, 305–334. doi:10.1037/0033-2909.129.2.305

- Lakin, J. L., Jefferis, V. E., Cheng, C. M., & Chartrand, T. L. (2003). The chameleon effect as social glue: Evidence for the evolutionary significance of nonconscious mimicry. *Journal of Nonverbal Behavior*, *27*, 145–162. doi:10.1023/A:1025389814290
- Lanzetta, J. T., & Englis, B. G. (1989). Expectations of cooperation and competition and their effects on observers' vicarious emotional responses. *Journal of Personality and Social Psychology*, *56*, 543–554. doi:10.1037/0022-3514.56.4.543
- Levenson, R. W., Oyama, O. N., & Meek, P. S. (1987). Greater reinforcement from alcohol for those at risk: Parental risk, personality risk, and sex. *Journal of Abnormal Psychology*, *96*, 242–253. doi:10.1037/0021-843X.96.3.242
- Maccoby, E. E. (1990). Gender and relationships: A developmental account. *American Psychologist*, *45*, 513–520. doi:10.1037/0003-066X.45.4.513
- Martin, C. S., & Sayette, M. A. (1993). Experimental design in alcohol administration research: Limitations and alternatives in the manipulation of dosage-set. *Journal of Studies on Alcohol and Drugs*, *54*, 750–761.
- McClure, E. B. (2000). A meta-analytic review of sex differences in facial expression processing and their development in infants, children, and adolescents. *Psychological Bulletin*, *126*, 424–453. doi:10.1037/0033-2909.126.3.424
- Mills, K. C., & Bisgrove, E. Z. (1983). Body sway and divided attention performance under the influence of alcohol: Dose-response differences between males and females. *Alcoholism: Clinical and Experimental Research*, *7*, 393–397. doi:10.1111/j.1530-0277.1983.tb05492.x

- Mooney, D. K., Fromme, K., Kivlahan, D. R., & Marlatt, G. A. (1987). Correlates of alcohol consumption: Sex, age, and expectancies relate differentially to quantity and frequency. *Addictive Behaviors, 12*, 235–240. doi:10.1016/0306-4603(87)90033-5
- Mulvihill, L. E., Skilling, T. A., & Vogel-Sprott, M. (1997). Alcohol and the ability to inhibit behavior in men and women. *Journal of Studies on Alcohol and Drugs, 58*, 600–605.
- Neff, L. A., & Karney, B. R. (2005). Gender differences in social support: A question of skill or responsiveness? *Journal of Personality and Social Psychology, 88*, 79–90.
doi:10.1037/0022-3514.88.1.79
- Niaura, R. S., Nathan, P. E., Frankenstein, W., Shapiro, A. P., & Brick, J. (1987). Gender differences in acute psychomotor, cognitive, and pharmacokinetic response to alcohol. *Addictive Behaviors, 12*, 345–356. doi:10.1016/0306-4603(87)90048-7
- Niedenthal, P. M. (2007). Embodying emotion. *Science, 316*, 1002–1005.
doi:10.1126/science.1136930
- Noldus Information Technology. (2010). *The Observer: Professional System for Collection, Analysis, and Presentation of Observational Data (Version XT)*. Wageningen, The Netherlands.
- Nolen-Hoeksema, S. (2004). Gender differences in risk factors and consequences for alcohol use and problems. *Clinical Psychology Review, 24*, 981–1010. doi:10.1016/j.cpr.2004.08.003
- Oberman, L. M., Winkielman, P., & Ramachandran, V. S. (2007). Face to face: Blocking facial mimicry can selectively impair recognition of emotional expressions. *Social Neuroscience, 2*, 167–178. doi:10.1080/17470910701391943
- Raudenbush, S. W., & Bryk, A. S. (2002). *Hierarchical linear models: Applications and data analysis methods* (Vol. 2). Thousand Oaks, CA: Sage Publications.

- Rohsenow, D. J. (1983). Drinking habits and expectancies about alcohol's effects for self versus others. *Journal of Consulting and Clinical Psychology, 51*, 752–756. doi:10.1037/0022-006X.51.5.752
- Rohsenow, D. J., & Marlatt, G. A. (1981). The balanced placebo design: Methodological considerations. *Addictive Behaviors, 6*, 107–122. doi:10.1016/0306-4603(78)90034-5
- Rosenbluth, J., Nathan, P. E., & Lawson, D. M. (1978). Environmental influences on drinking by college students in a college pub: Behavioral observation in the natural environment. *Addictive Behaviors, 3*, 117–121.
- Ruch, W. (1994). Extraversion, alcohol, and enjoyment. *Personality and Individual Differences, 16*, 89–102. doi:10.1016/0191-8869(94)90113-9
- Sastry, N. (1997). A nested frailty model for survival data, with an application to the study of child survival in northeast Brazil. *Journal of the American Statistical Association, 92*, 426–435. doi:10.1080/01621459.1997.10473994
- Sayette, M. A., Breslin, F. C., Wilson, G. T., & Rosenblum, G. D. (1994). Parental history of alcohol abuse and the effects of alcohol and expectations of intoxication on social stress. *Journal of Studies on Alcohol and Drugs, 55*, 214–223.
- Sayette, M. A., Creswell, K. G., Dimoff, J. D., Fairbairn, C. E., Cohn, J. F., Heckman, B. W., ... Moreland, R. L. (2012). Alcohol and group formation: A multimodal investigation of the effects of alcohol on emotion and social bonding. *Psychological Science, 23*, 869–878. doi:10.1177/0956797611435134
- Sayette, M. A., Dimoff, J. D., Levine, J. M., Moreland, R. L., & Votruba-Drzal, E. (2012). The effects of alcohol and dosage-set on risk-seeking behavior in groups and individuals. *Psychology of Addictive Behaviors, 26*, 194–200. doi:10.1037/a0023903

- Sayette, M. A., Martin, C. S., Perrott, M. A., Wertz, J. M., & Hufford, M. R. (2001). A test of the appraisal-disruption model of alcohol and stress. *Journal of Studies on Alcohol*, *62*, 247–256.
- Scheel, I., Aldrin, M., Frigessi, A., & Jansen, P. A. (2007). A stochastic model for infectious salmon anemia (ISA) in Atlantic salmon farming. *Journal of The Royal Society Interface*, *4*, 699–706. doi:10.1098/rsif.2007.0217
- Senchak, M., Leonard, K. E., & Greene, B. W. (1998). Alcohol use among college students as a function of their typical social drinking context. *Psychology of Addictive Behaviors*, *12*, 62–70. doi:10.1037/0893-164X.12.1.62
- Sher, K. J. (1987). Stress Response Dampening. In H. T. Blane & K. E. Leonard (Eds.), *Psychological theories of drinking and alcoholism*. New York: Guilford Press.
- Sher, K. J., Grekin, E. R., & Williams, N. A. (2005). The development of alcohol use disorders. *Annual Review of Clinical Psychology*, *1*, 493–523.
doi:10.1146/annurev.clinpsy.1.102803.144107
- Single, E., & Wortley, S. (1993). Drinking in various settings as it relates to demographic variables and level of consumption: Findings from a national survey in Canada. *Journal of Studies on Alcohol and Drugs*, *54*, 590–599.
- Sonnby-Borgstrom, M., Jönsson, P., & Svensson, O. (2008). Gender differences in facial imitation and verbally reported emotional contagion from spontaneous to emotionally regulated processing levels. *Scandinavian Journal of Psychology*, *49*, 111–122.
doi:10.1111/j.1467-9450.2008.00626.x

- Steele, C. M., & Josephs, R. A. (1988). Drinking your troubles away II: An attention-allocation model of alcohol's effect on psychological stress. *Journal of Abnormal Psychology, 97*, 196–205. doi:10.1037/0021-843X.97.2.196
- Steele, C. M., & Josephs, R. A. (1990). Alcohol myopia: Its prized and dangerous effects. *American Psychologist, 45*, 921–933. doi:10.1037/0003-066X.45.8.921
- Stel, M., & van Knippenberg, A. (2008). The role of facial mimicry in the recognition of affect. *Psychological Science, 19*, 984–985. doi:10.1111/j.1467-9280.2008.02188.x
- Stel, M., & Vonk, R. (2010). Mimicry in social interaction: Benefits for mimickers, mimicked, and their interaction. *British Journal of Psychology, 101*, 311–323.
- Stoolmiller, M., & Snyder, J. (2006). Modeling heterogeneity in social interaction processes using multilevel survival analysis. *Psychological Methods, 11*, 164–177. doi:10.1037/1082-989X.11.2.164
- Substance Abuse and Mental Health Services Administration. (2012). National Survey on Drug Use and Health: Summary of National Findings. Retrieved from <http://www.samhsa.gov>
- Sutker, P. B., Allain, A. N., Brantley, P. J., & Randall, C. L. (1982). Acute alcohol intoxication, negative affect, and autonomic arousal in women and men. *Addictive Behaviors, 7*, 17–25. doi:10.1016/0306-4603(82)90020-X
- Testa, M., Fillmore, M. T., Norris, J., Abbey, A., Curtin, J. J., Leonard, K. E., ... George, W. H. (2006). Understanding alcohol expectancy effects: Revisiting the placebo condition. *Alcoholism: Clinical and Experimental Research, 30*, 339–348.
- Therneau, T. (2011). Coxme: Mixed effects Cox models. *R Package Version*.

- Tiedens, L. Z., & Fragale, A. R. (2003). Power moves: Complementarity in dominant and submissive nonverbal behavior. *Journal of Personality and Social Psychology*, *84*, 558–568. doi:10.1037/0022-3514.84.3.558
- Van Leeuwen, M. L., van Baaren, R. B., Martin, D., Dijksterhuis, A., & Bekkering, H. (2009). Executive functioning and imitation: Increasing working memory load facilitates behavioural imitation. *Neuropsychologia*, *47*, 3265–3270. doi:10.1016/j.neuropsychologia.2009.06.005
- Wilson, G. T. (1988). Alcohol and anxiety. *Behaviour Research and Therapy*, *26*, 369–381. doi:10.1016/0005-7967(88)90070-8
- Wood, W., & Eagly, A. H. (2010). Gender. In S. T. Fiske, D. T. Gilbert, & G. Lindzey (Eds.), *Handbook of social psychology* (5th ed., Vol. 1, pp. 629–667). New York: John Wiley.

Table 1. Beverage Manipulation Check

	Alcohol		Placebo		Control		<i>F</i>
	Mean	SD	Mean	SD	Mean	SD	
BAC after drinking	0.055 ^a	0.012	0.001 ^b	0.001	0.001 ^b	0.001	4825.72**
BAC 40-min after drinking†	0.062 ^a	0.011	0.001 ^b	0.001	—	—	7116.15**
SIS after drinking	38.50 ^a	17.31	14.90 ^b	10.44	0.20 ^c	1.49	647.70**
SIS 40-min after drinking†	35.12 ^a	16.90	8.90 ^b	10.80	—	—	410.12**
Highest Intox.	43.53 ^a	18.71	16.15 ^b	11.11	0.61 ^c	3.19	698.07**
Vodka Estimate (in ounces)	7.11 ^a	9.85	4.64 ^b	5.44	0.05 ^c	0.43	70.80**

* $p < .05$ ** $p < .001$

† control participants not asked to provide these data


Notes. BAC = blood alcohol concentration. SIS = subjective intoxication scale. SIS and Highest Intox. were scored on scales ranging from 0 to 100. Groups with non-overlapping superscripts differed significantly ($p < .05$).

Table 2. Percentage of smiles leading to a mutual smile

Group Gender Makeup	Alcohol % (N)	Placebo % (N)	Control % (N)	Total % (N)
All Males	50.4% (614)	37.4% (321)	38.2% (434)	42.6% (1369)
2 Males 1 Female	48.8% (780)	44.4% (490)	45.7% (651)	46.5% (1921)
1 Male 2 Female	49.8% (714)	49.1% (683)	45.7% (558)	48.4% (1955)
All Females	49.4% (822)	48.2% (679)	47.9% (659)	48.6% (2160)
Total	49.6% (2930)	45.6% (2173)	44.6% (2302)	46.8% (7405)


Percentage of smiles initiated within groups that led to a smile that was shared with another group member (versus end with eliciting a responding smile). Note that percentages are not intended to sum to 100%, since non-reciprocated smiles are not represented within the table.

Figure 1. Cumulative hazard of a smile developing into a mutual smile across group gender composition



Smiles that ended without eliciting a responding smile are represented by + symbol. Lines become thinner across horizontal axis as observations become more sparse (the majority of smiles either developed into a mutual smile or ended un-reciprocated within 5 seconds of their initiation.)

Figure 2. Cumulative hazard of a smile developing into a mutual smile according to drink condition and the gender of the smile initiator's group-mates



Smiles that ended without eliciting a responding smile are represented by + symbol. The gender of the smile initiator him/herself did not significantly impact mutual smile hazard in any drink condition